

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

DEMOGRAPHICS US Census: Official Population Estimates	DISTRICT								COMPARISONS	
	Aroostook	Central	Cumber.	Downeast	Mid Coast	Penquis	Western	York	MAINE STATE	UNITED STATES
Total Population [2008]	71,676	172,336	276,047	85,636	149,988	165,612	193,475	201,686	1,316,456	304,059,724
Total Percent of Maine Population [2008]	5.4	13.1	21.0	6.5	11.4	12.6	14.7	15.3	100	n/a
Population Density, people per square mile [2008]	10.7	35.9	330.4	20.6	83.1	22.5	45.6	203.6	42.7	86.1
Selected Ages (US Census Official Population Estimates) [2008]										
<5 Years of Age, percent and count [2008]	4.9 3,527	5.3 9,110	5.6 15,505	5.2 4,425	5.2 7,799	5.5 9,096	5.8 11,271	5.3 10,726	5.4 71,459	6.9 20,480,587
<18 Years of Age, percent and count [2008]	19.6 14,029	20.7 35,722	21.5 59,301	19.6 16,805	20.4 30,660	20.2 33,438	21.4 41,369	21.6 43,543	20.9 274,867	24.3 73,941,848
65 Years & Older, percent and count [2005-2007]	17.6 12,669	14.6 25,100	13.5 37,029	16.8 14,408	16.1 24,107	13.8 22,862	14.4 27,939	13.6 25,429	14.4 183,402	12.5 37,265,110
Race/Ethnicity Counts (one race alone or in combination with one or more other races, except where noted.** US Census Official Population Estimates)										
White, percent and count [2008]	97.2 69,653	97.2 169,464	95.6 263,911	97.0 83,057	98.5 147,758	97.4 161,345	97.8 189,213	98.1 197,932	97.4 128,2333	81.3 247,112,954
Black, percent and count, [2008]	0.8 608	0.8 1,447	2.5 6,960	0.7 593	0.9 1,297	1.0 1,753	1.6 3,055	1.0 2,064	1.4 17,777	13.5 41,126,808
American Indian & Alaskan Native, percent and count [2008]	2.0 1,429	1.0 1,717	0.8 2,323	2.6 2,240	0.9 1,401	1.4 2,308	0.9 1,702	0.6 1,290	1.1 14,410	0.8 2,328,982
Asian, percent and count [2008]	0.8 605	0.9 1,527	2.2 5,958	0.9 738	0.7 1,033	1.1 1,762	0.8 1,617	1.1 2,181	1.8 15,421	4.4 13,237,698
Native Hawaiian and Other P.I., percent and count [2008]	0.03 25	0.05 91	0.14 398	0.04 30	0.04 61	0.08 140	0.06 114	0.05 91	0.07 950	0.14 434,561
Hispanic of any race**, percent and count [2008]	1.1 790	1.1 1,826	1.8 5,028	1.2 1,023	1.1 1,629	1.0 1,659	1.2 2,403	1.2 2,456	1.3 16,814	15.4 46,943,613
Non-Hispanic**, percent and count [2008]	98.9 70,886	98.9 170,510	98.2 271,019	98.8 84,613	98.9 148,359	99.0 163,953	98.8 191,072	98.8 199,230	98.7 1,299,642	84.66 257,116,111
Franco-American, count and percent where applicable [2000]	25,374	46,580	47,185	9,083	19,132	46,580	58,457	52,868	305,259 (22%)	0.8%

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

DEMOGRAPHICS (cont.)	DISTRICT								COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES
Children Eligible for Free or Reduced Lunch Program, percent and count [FY2009]	49.4 (48.5- 50.4) 5,449	44.2 (43.7- 44.8) 11,931	27.3 (26.9- 27.9) 11,268	44.1 (43.2- 45.0) 5,135	41.6 (40.9- 42.3) 8,158	43.5 (42.9- 44.2) 10,769	47.9 (47.3- 48.5) 14,051	30.2 (29.6- 30.7) 8,547	39.1 (38.8- 39.3) 75,358	n/a
Adults with Lifetime Educational Attainment Less Than High School, percent and count [2000]	23.1 11,882	16.1 18,418	9.9 17,946	15.3 9,164	13.0 13,139	14.9 16,068	18.6 23,577	13.5 17,225	14.6 127,004	9.4
Single-Parent Households with Children <18 years, percent and count [2000]	7.7 2,323	10.4 7,077	8.4 9,117	8.8 3,157	9.0 5,372	9.5 6,197	10.4 7,887	9.1 6,788	9.2 47,918	7.2
Householders ≥ 65 Living Alone, percent and count [2000]	13.1 3,977	10.5 7,145	10.2 11,015	12.2 4,386	11.0 6,549	10.4 6,829	10.9 8,317	9.7 7,233	10.7 55,447	9.2
People Who Speak a Language Other Than English >5 yrs old, percent and count [2000]	24.1 16,880	6.8 10,735	5.9 14,888	4.3 3,498	3.5 4,798	4.4 6,818	11.1 19,771	9.4 16,578	7.8 93,966	17.9
Disability among those >5 yrs old, percent and count [2000]	25.2 17,438	21.4 33,513	17.1 42,102	22.2 17,830	18.8 25,250	20.3 30,721	21.8 38,143	18.9 32,913	20.0 237,910	19.3
Adults with Disability, percent [2008]	25.0 (±5.4)	27.1 (±3.7)	21.8 (±2.9)	23.0 (±3.7)	24.1 (±2.8)	24.3 (±3.6)	25.5 (±3.3)	22.6 (±3.7)	24.2 (±1.2)	n/a
Individuals Living in Poverty, percent and count [2007]	17.4 12,273	13.9 24,020	9.7 25,837	13.5 11,574	11.1 16,651	13.3 21,986	14.0 27,150	8.2 16,271	12.2 155,764	13.0 38,052,247

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

MATERNAL/CHILD HEALTH INDICATORS	DISTRICT									COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Infant Mortality, rate per 1,000 live births [2003-2007]	6.6 (4.2-9.9)	5.7 (4.4-7.5)	5.7 (4.6-7.0)	4.4 (2.6-6.8)	5.1 (3.6-6.9)	6.3 (4.7-8.1)	6.8 (5.4-8.6)	5.7 (4.3-7.5)	6.0 (5.5-6.6)	6.8 [2007]	WA 4.7 [2006]
Live Births with Low Birth Weight <2500 grams, percent of live births [2008]	5.9 (4.3-7.9)	6.0 (4.9-7.3)	6.8 (5.9-7.8)	5.7 (4.2-7.6)	6.6 (5.4-8.1)	7.4 (6.1-8.8)	7.0 (6.0-8.2)	7.4 (6.3-8.7)	6.7 (6.3-7.1)	8.2 [2007]	AK 5.7 [2007]
Infants Born to Women Receiving First Trimester Prenatal Care, percent [2008]	87.7 (81.1-94.7)	83.1 (78.8-87.5)	87.5 (84.1-91.0)	83.6 (77.5-90.0)	89.6 (84.7-94.6)	84.3 (80.0-88.8)	88.6 (84.7-92.6)	88.4 (84.3-92.6)	86.8 (85.3-88.4)	83.2 [2006]	MA 88.6 [2006]
Teen Births Ages 15-19 years, rate per 1,000 female population [2006-2008]	28.8 (25.0- 33.1)	33.7 (30.9- 36.6)	17.0 (15.5- 18.6)	31.0 (27.2- 35.2)	29.3 (26.5- 32.4)	24.6 (22.3- 26.9)	34.4 (31.8- 37.1)	19.8 (17.9- 21.9)	26.0 (25.2- 26.9)	42.5 [2007]	VT 10.4 [2006]
Pregnancy Rate 15-44 Females, rate per 1,000 female population [2006-2008]	59.5 (57.1-62.0)	61.9 (60.4-63.5)	70.2 (68.9-71.5)	65.3 (63.0-67.7)	64.9 (63.2-66.7)	59.7 (58.2-61.2)	67.6 (66.1-69.1)	61.8 (60.4-63.2)	64.6 (64.0-65.2)	103.2 [2005]	n/a
Preterm Births, <37 weeks, percent of live births [2008]	7.2 (5.4-9.4)	8.5 (7.2-10.0)	9.3 (8.2-10.4)	5.7 (4.2-7.6)	8.8 (7.4-10.5)	9.2 (7.8-10.7)	8.2 (7.1-9.5)	9.7 (8.4-11.1)	8.7 (8.2-9.2)	n/a	n/a
C-Section Births, percent of live births [2008]	31.0 (27.2-35.3)	31.2 (28.6-34.0)	30.9 (28.9-33.0)	28.3 (24.8-32.1)	29.5 (26.8-32.5)	28.9 (26.4-31.6)	33.3 (31.0-35.8)	29.3 (27.0-31.8)	30.6 (29.6-31.5)	31.1 [2007]	AK 22.6 [2007]
Infants Born to Women who Used Tobacco During Last 3 Months of Pregnancy, percent of live births [2004-2007]	26.1 (20.1-33.1)	18.8 (15.4-22.6)	10.9 (8.8-13.5)	21.8 (16.6-28.0)	18.2 (14.7-22.3)	26.2 (22.1-30.6)	22.9 (19.4-26.8)	14.9 (11.8-18.6)	18.6 (17.3-20.0)	n/a	NJ 5.7 [2006]
Children with Special Health Needs, percent [2005-2006]	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	17.7 (16.5-18.8)	13.9 (13.7-14.1 [2005-06])	CA 9.9 [2005-06]

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

MATERNAL/CHILD HEALTH INDICATORS Continued	DISTRICT									COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Children 6-17 years in Special Education, percent	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	14.9 [2007-2008]	11.4 [2007-2008]	n/a
Percent of Children 0-19 Years Enrolled in MaineCare, percent and count [2008]	49.2 (48.5- 49.9) 9,2140	46.0 (45.5- 46.4) 20,762	30.0 (29.6- 30.3) 20,691	45.7 (45.1- 46.4) 9,822	41.0 (40.5- 41.5) 15,348	44.7 (44.2- 45.2) 19,143	50.4 (50.0- 50.9) 25,434	32.3 (31.9- 32.7) 16,241	41.5 (41.4- 41.7) 139,378	n/a	n/a
Breastfeeding Initiation, percent [2004-2007]	71.1 (64.0- 77.3)	80.8 (76.9- 84.1)	84.6 (81.8- 87.1)	77.6 (71.3- 82.9)	78.9 (74.7- 82.7)	73.9 (69.5- 77.9)	79.4 (75.6- 82.8)	75.4 (71.1- 79.2)	78.9 (77.5- 80.3)	n/a	OR 93.7 [2006]
HEALTH & WELLNESS INDICATORS											
Adolescent Smoking Prevalence, 6-12 Graders, percent [2008]	14.5 (12.7-15.7)	11.7 (11.2-12.2)	11.3 (10.7-11.8)	14.1 (12.6-15.6)	13.0 (11.9-14.0)	12.6 (11.4-13.8)	11.0 (10.1-11.8)	12.1 (10.4-13.8)	12.1 (11.7-12.5)	n/a	n/a
Adult Current Smoking Prevalence, percent [2008]	21.9 (±5.7)	20.7 (±4.1)	13.3 (±2.9)	17.8 (±3.8)	18.4 (±3.1)	19.9 (±3.6)	21.3 (±3.3)	15.4 (±3.6)	18.2 (±1.3)	18.4	CA 14.0 [2008]
High School Youth Overweight or Obese, percent [2007]	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	25.9 (22.3, 29.8)	28.8	n/a
Adults Overweight or Obese, percent [2008]	69.1 (±6.2)	58.7 (±4.7)	56.5 (±3.9)	61.5 (±5.1)	58.8 (±3.6)	68.4 (±4.2)	65.9 (±3.9)	62.3 (±4.6)	61.8 (±1.6)	63.2	CO 55.3 [2008]
Adults Reporting Fair or Poor Health Status in last 30 days, percent [2008]	17.8 (±4.5)	13.5 (±2.8)	10.9 (±2.0)	13.2 (±2.9)	10.9 (±2.1)	14.4 (±2.8)	14.6 (±2.8)	11.6 (±3.0)	13.0 (±1.0)	14.4	UT 10.6 [2008]
Adults with >= 14 Unhealthy Days per Month, percent [2008]	23.1 (±5.9)	19.6 (±3.4)	16.3 (±3.0)	16.0 (±3.4)	15.7 (±2.6)	17.7 (±3.1)	16.9 (±2.8)	18.6 (±3.6)	17.7 (±1.2)	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Overall Cancer Incidence, age-adjusted rate per 100,000 [2004-2006]	508.6 (482.5-535.8)	527.3 (509.1-546.0)	513.2 (498.5-528.1)	574.3 (548.7-600.8)	527.6 (508.7-547.1)	581.1 (561.1-601.6)	526.2 (508.9-544)	525.1 (508.0-542.7)	533 (526.3-539.7)	456.2 (453.7-458.7)	n/a
Overall Cancer Mortality, age-adjusted rate per 100,000 [2003-2007]	197.4 (±12.4)	200.6 (±8.7)	194.9 (±6.9)	210.8 (±11.8)	187.7 (±8.6)	210.8 (±9.3)	203.6 (±8.2)	190.0 (±7.9)	198.5 (±3.1)	180.7 [2005]	UT 135.6 [2006]
Lung Cancer Incidence, age-adjusted rate per 100,000 [2004-2006]	88.1 (77.7- 99.6)	84 (76.8- 91.6)	70.9 (65.5- 76.6)	96.9 (86.8- 108)	75.3 (68.4- 82.8)	90.9 (83.2- 99.2)	79.5 (72.9- 86.6)	73.7 (67.4- 80.5)	80.3 (77.7- 82.9)	60.0 (59.1-61.0) [2006]	n/a
Lung Cancer Mortality, age-adjusted rate per 100,000 [2003-2007]	62.9 (±7.0)	64.6 (±4.9)	56.7 (±3.8)	59.7 (±6.3)	56.3 (±4.7)	63.1 (±5.1)	63.2 (±4.6)	56.1 (±4.3)	59.9 (±1.7)	51.5 [2006]	UT 23.3 [2006]
Colorectal Cancer Incidence, age-adjusted rate per 100,000 [2004-2006]	58.3 (49.8- 67.9)	52.1 (46.5- 58.2)	48.2 (43.8- 52.9)	57.8 (50- 66.5)	51.3 (45.6- 57.6)	65.4 (58.8- 72.5)	55.2 (49.7- 61.1)	52.2 (46.9- 57.9)	54.2 (52.1- 56.4)	45.4 (44.7-46.2) [2006]	n/a
Colorectal Cancer Mortality, age-adjusted rate per 100,000 [2003-2007]	20.7 (±4.0)	17.5 (±2.6)	19.5 (±2.2)	17.9 (±3.4)	14.8 (±2.4)	19.4 (±2.8)	18.5 (±2.5)	17.2 (±2.4)	18.1 (±0.9)	17.1 [2006]	UT 13.0 [2006]
Sigmoidoscopy or Colonoscopy Screening Ever Had by Adults Age 50 and Older, percent [2008]	67.6 (±6.7)	75.1 (±4.2)	76.5 (±3.6)	63.9 (±5.2)	70.5 (±3.3)	69.0 (±4.7)	72.9 (±4.0)	77.8 (±4.2)	72.6 (±1.5)	62.2	LA 52.6 [2008]
Female Breast Cancer Incidence, age-adjusted rate per 100,000 [2004-2006]	106.4 (90.1- 125)	130.0 (117.8-143.1)	126.1 (116.5-136.5)	118.4 (102.8-136.0)	136.9 (123.8-151.1)	134.2 (121.2-148.2)	126.6 (115.0-139.1)	132.3 (120.8-144.7)	128.1 (123.6-132.6)	123 (121.3-124.8) [2006]	n/a
Female Breast Cancer Mortality, age-adjusted rate per 100,000 [2003-2007]	18.6 (±5.9)	19.1 (±3.7)	22.7 (±3.1)	25.1 (±6.5)	19.9 (±3.9)	25.9 (±4.4)	25.8 (±4.0)	21.9 (±3.7)	22.5 (±1.4)	23.5 [2006]	MT 18.8 [2006]

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS Continued	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Mammogram in Past Two Years Among Women 40 and Older, percent [2008]	85.0 (±5.8)	82.7 (±4.4)	84.3 (±3.5)	77.6 (±5.6)	82.6 (±3.2)	84.8 (±4.2)	83.0 (±3.7)	85.1 (±4.2)	83.3 (±1.5)	76	MA 84.9 [2008]
Pap Smear in Past 3 Years Among Women >=18 years, percent [2008]	90.0 (±5.3)	82.8 (±5.4)	88.4 (±3.7)	83.6 (±5.3)	84.4 (±4.5)	88.3 (±4.5)	85.9 (±4.6)	88.0 (±5.5)	86.3 (±1.7)	82.9	MA 87.6 [2008]
Prostate Cancer Incidence age-adjusted rate per 100,000 [2004-2006]	128.5 (110.0-149.5)	159.6 (145.0-175.2)	167.1 (154.7-180.2)	184.9 (164.3-207.6)	168 (152.6-184.6)	176 (160-193.2)	165.6 (151.4-180.8)	156.8 (143.2-171.3)	164.6 (159.2-170.2)	163.1 (160.8-165.3)	n/a
Prostate Cancer Mortality age-adjusted rate per 100,000 [2003-2007]	11.2 (±5.6)	27.2 (±5.3)	27.7 (±4.3)	25.2 (±7.3)	29.3 (±5.4)	25.1 (±6.0)	21.9 (±4.8)	25.1 (±4.7)	25.0 (±1.8)	23.5 [2006]	HI 13.5 [2006]
Major CVD Deaths, age-adjusted rate per 100,000 and counts [2007]	258.1 (±30.5) 271	245.6 (±20.9) 522	184.5 (±14.6) 618	256.5 (±28.1) 312	235.5 (±21.0) 475	262.2 (±22.9) 491	224.7 (±18.9) 545	206.3 (±17.9) 502	226.4 (±7.2) 3736	261.2 [2006]	MN 190.1 [2006]
Stroke Hospitalizations, age-adjusted rate per 10,000 [2008]	23.8 (20.9- 27.0)	17.5 (15.7- 19.4)	16.7 (15.3- 18.2)	21.2 (18.7- 24.0)	16.5 (14.7- 18.4)	22.4 (20.4- 24.7)	18.3 (16.6- 20.1)	16.0 (14.4- 17.7)	18.3 (17.7- 19.0)	n/a	n/a
Acute Myocardial Infarction Hospitalizations, age-adjusted rate per 10,000 [2008]	46.1 (41.8- 50.7)	32.9 (30.5- 35.4)	16.2 (14.8- 17.6)	38.1 (34.6- 41.8)	23.0 (20.9- 25.2)	33.2 (30.7- 36.0)	22.4 (20.5- 24.4)	22.0 (20.2- 24.0)	26.3 (25.5- 27.1)	n/a	n/a
High Blood Pressure Among Adults, percent [2008]	31.5 (±5.3)	31.6 (±3.7)	27.5 (±3.1)	32.7 (±4.3)	29.3 (±2.7)	33.1 (±3.8)	33.5 (±3.4)	30.0 (±3.5)	30.9 (±1.3)	n/a	n/a
High Cholesterol Among Adults, percent [2008]	44.1 (±6.0)	40.8 (±3.9)	37.2 (±3.4)	45.7 (±4.9)	39.9 (±3.2)	44.2 (±4.1)	44.4 (±3.8)	35.5 (±4.0)	40.7 (±1.4)	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS Continued	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Diabetes Prevalence Among Adults, percent [2008]	9.8 (±3.2)	9.8 (±2.3)	8.8 (±1.9)	8.1 (±2.2)	6.6 (±1.3)	10.2 (±2.4)	6.6 (±1.7)	6.9 (±2.0)	8.3 (±0.7)	8.3	MN 5.9 [2008]
Diabetes Hospitalizations, age-adjusted rate per 10,000 [2008]	13.5 (11.0- 16.4)	10.7 (9.3- 12.3)	9.5 (8.4- 10.8)	13.3 (11.0- 15.9)	9.8 (8.3- 11.4)	16.9 (15.0-18.9)	12.4 (10.9- 14.0)	11.1 (9.7- 12.7)	11.8 (11.2- 12.4)	n/a	n/a
Adults with Diabetes Who Have Received a Hemoglobin A1c Test at Least Once Yearly, percent [2008]	89.9 (±8.8)	91.1 (±8.1)	93.7 (±6.9)	94.7 (±4.3)	95.7 (±3.2)	96.6 (±2.9)	89.0 (±6.3)	93.9 (±4.6)	92.9 (±2.3)	n/a	n/a
Adults With Diabetes Who Have Taken a Diabetes Management Course, percent [2008]	70.1 (±8.0)	70.7 (± 6.5)	62.7 (±7.0)	52.2 (±9.6)	62.8 (±6.1)	59.0 (±7.4)	51.6 (±7.9)	56.5 (±8.2)	60.5 (±5.4)	n/a	n/a
Diabetes Deaths, age-adjusted rate per 100,000 [2003-2007]	25.7 (±4.5)	27.1 (±3.2)	18.9 (±2.2)	31.0 (±4.5)	17.5 (±2.6)	26.3 (±3.3)	26.9 (±3.0)	21.9 (±2.7)	23.6 (±1.1)	23.2 [2006]	NV 12.8 [2006]
Adults with Asthma, percent [2008]	10.6 (±4.2)	11.3 (±2.8)	9.3 (±2.0)	10.3 (±3.2)	7.6 (±1.7)	12.6 (±3.3)	11.6 (±2.6)	9.4 (±2.9)	10.3 (±1.0)	8.8	FL 6.6 [2008]
Child and Youth Asthma, <18 Years Old, percent [2007]	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	9.0 (7.4-10.6)	9.0 (8.6-9.5)	n/a
Asthma Emergency Department Visits, age-adjusted rate per 10,000 [2007]	92.5 (85.2- 100.2)	88.8 (84.2- 93.5)	52.5 (49.7- 55.4)	73.3 (67.3- 79.6)	57.5 (53.5- 61.8)	60.1 (56.3- 64.0)	76.0 (72.0- 80.2)	53.3 (50.0- 56.8)	65.9 (64.5- 67.4)	n/a	n/a
Chronic Lower Respiratory Disease Deaths, 45 Years and Older, rate per 100,000 [2003-2007]	142.4 (±17.7)	157.1 (±13.1)	127.5 (±9.7)	144.6 (±16.6)	146.6 (±13.0)	143.0 (±13.1)	173.9. (±13.0)	121.0 (±10.9)	143.7 (±4.5)	114.6 [2006]	HI 55.3 [2006]

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

CHRONIC DISEASE INDICATORS Continued	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Adults With a Routine Dental Visit in Past Year, percent [2008]	67.6 (±6.1)	65.3 (±4.4)	76.8 (±3.5)	67.9 (±4.5)	72.0 (±3.3)	68.8 (±4.0)	65.0 (±3.8)	74.5 (±4.3)	70.2 (±1.5)	n/a	n/a
Adults Who Have Lost Some Teeth Due to Gum Disease or Tooth Decay, percent [2008]	45.5 (±6.5)	44.4 (±4.4)	33.9 (±3.4)	42.2 (±4.8)	40.6 (±3.4)	44.4 (±4.4)	43.5 (±3.9)	38.8 (±4.2)	41.0 (±1.5)	n/a	n/a
Adults Who Have Lost All Teeth Due to Gum Disease or Tooth Decay, percent [2008]	11.1 (±3.1)	7.0 (±1.6)	5.5 (±1.4)	6.3 (±1.9)	5.5 (±1.2)	9.8 (±2.1)	8.7 (±1.7)	5.7 (±1.7)	7.2 (±0.6)	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ENVIRONMENTAL HEALTH INDICATORS	DISTRICT								COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES
Emergency Department Visits for Carbon Monoxide Poisoning, rate per 100,000 [2002-2006]	8.2 (±3.1)	10.5 (±2.3)	5.9 (±1.4)	7.7 (±2.8)	6.7 (±1.9)	7.3 (±1.9)	11.5 (±2.2)	7.9 (±1.8)	8.1 (±0.7)	n/a
Carbon Monoxide Detector in the Home, percent [2004]	29.5 (±7.0)	35.0 (±4.8)	35.6 (±4.1)	41.7 (±6.1)	33.7 (±3.7)	37.9 (±5.2)	33.9 (±4.6)	36.6 (±4.9)	35.5 (±1.7)	n/a
Lead Screening Among 1-Year Old Children, percent [2003-2007]	67.1 (±1.6)	33.1 (±1.0)	45.2 (±0.8)	61.0 (±1.5)	43.1 (±1.1)	54.6 (±1.1)	51.9 (±1.0)	50.9 (±0.9)	48.7 (±0.4)	n/a
Elevated Blood Lead Levels Among Screened 1-Year Old Children, percent [2003-2007]	0.4 (±0.3)	1.6 (±0.5)	1.5 (±0.3)	1.2 (±0.5)	2.0 (±0.5)	1.4 (±0.3)	1.5 (±0.3)	1.3 (±0.3)	1.4 (±0.1)	1.58 (<72 mos old) [2005]
District Community Water Systems Meeting All Health Based Standards, percent [2008]	78	84	84	84	83	94	84	93	86	89.3
District Population Served by Community Water Meeting All Health-Based Standards, percent [2008]	66	20	98	82	76	77	96	95	82	n/a
District Population Served by Community Water Systems with Source Water Protection in Place, percent [2008]	71	92	93	77	92	85	96	100	92	89.4
Total Number of Licensed Site Evaluators That Design Septic Systems to Insure Proper Wastewater Disposal [2009]	6	63	51	19	40	48	31	27	285	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ENVIRONMENTAL HEALTH INDICATORS Continued	DISTRICT								COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES
Total Number of Voluntary Certified Inspectors That Perform On-Site Septic System Inspections to Insure Proper Wastewater Disposal [2009]	22	89	89	48	54	48	73	54	477	n/a
Total Number of Subsurface Wastewater System (Septic System) Permits Issued [2006]	557	1309	1512	1045	1502	958	1546	1178	9607	n/a
Total Number of Eating and Lodging Licenses Issued Per Year [2009]	268	835	1245	597	769	631	766	1034	6145	n/a
Total Number of Body Artist Licenses Issued Per Year [2009]	11	44	78	4	4	57	18	21	237	n/a
Total Number of Eating and Lodging Place Inspections Per Year [2008]	315	492	426	150	471	438	588	438	3318	n/a
Total Number of Body Artist Inspections Per Year [2008]	6	10	5	3	2	17	7	7	57	n/a
Total Number of Failed Health Inspections Per Year [2008]	11	5	11	2	7	8	14	5	63	n/a
Total Number of Voluntary Closures for Failed Inspections Per Yr [2008]	3	5	1	1	1	6	0	0	17	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ENVIRONMENTAL HEALTH INDICATORS Continued	DISTRICT								COMPARISONS	
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES
Homes with Radon in Air Concentrations Above the 4pCi/l "Take Action to Reduce" Level, percent [1989]	43.1 (±4.9)	25.0 (±4.5)	40.9 (±4.3)	23.7 (±4.4)	19.3 (±3.8)	19.8 (±3.6)	34.2 (±4.5)	41.8 (±5.5)	30.9 (±1.6)	n/a
Homes That Have Tested Indoor Air for Radon, percent [2006]	19.5 (±5.9)	25.1 (±4.3)	40.7 (±4.1)	22.0 (±4.7)	27.4 (±3.5)	23.5 (±4.2)	33.9 (±4.6)	30.0 (±4.6)	28.0 (±1.6)	n/a
Total Number of Licensed Radon Analysis Labs, Testers, and Mitigators That Identify and Reduce Elevated Radon Concentrations in the Air or Water of Buildings [2009]	41	50	59	46	51	43	65	53	110	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

INFECTIOUS DISEASE INDICATORS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Kindergarteners Exempted from Childhood Vaccine for Philosophical Reasons, count [2007]	7	16	85	45	41	39	42	34	316		
Children Immunized with the 4:3:1:3:3:1 Vaccination Series by 24 Months of Age, percent [2004-2005] <small>(4:3:1:3:3:1 means at least 4 DTP/DT/DTaP, 3 Polio, 1 Measles-containing, 3 Hib, 3 Hepatitis B, and 1 Varicella vaccine)</small>	n/a	n/a	75.0 (±6.3)	n/a	n/a	n/a	n/a	77.4 (±6.5)	67.1 (±6.2) [2008]	71.3 (±1.2) [2008]	NH 79.6 [2008]
Influenza Vaccine Past Year for Adults >65 years, percent [2008]	68.1 (±9.9)	78.4 (±5.9)	75.6 (±5.6)	76.8 (±6.9)	75.9 (±4.3)	78.5 (±6.2)	69.1 (±5.9)	74.7 (±6.6)	74.6 (±2.2)	71.1	NH 78.1 [2008]
Ever Had Pneumococcal Vaccine, >=65 Years, percent [2008]	68.2 (±7.6)	72.8 (±4.8)	75.6 (±4.3)	73.1 (±5.2)	66.9 (±3.7)	82.0 (±4.1)	70.8 (±4.5)	64.1 (±5.3)	71.7 (±1.7)	n/a	n/a
Reported Cases of Chlamydia, count [2008]	64	424	644	127	250	342	465	278	2,604	n/a	n/a
Reported Cases of Gonorrhea, average annual count [2004-2008]	5	12	51	6	8	12	31	17	141	n/a	n/a
Reported Cases of Chronic Hepatitis C, average annual count [2006-2008]	39	148	400	50	111	167	152	155	1,344	n/a	n/a
Late Diagnosis of HIV/AIDS, diagnosis within 12 mos. of 1st HIV diagnosis, 5-yr count [2004-2008]	2	13	34	6	10	6	19	21	111	n/a	n/a
Reported Cases of Lyme Disease, crude rate per 100,000 [2008]	5.6	71.4	82.6	18.7	114.0	8.5	31.5	144.3	69.0	11.6	n/a
Reported Cases of Salmonella, crude rate per 100,000 [2008]	9.8	16.2	14.1	10.5	14	7.2	7.8	13.9	12.1	16.8	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

INJURY AND VIOLENCE INDICATORS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Motor Vehicle Traffic Crash Deaths, age-adjusted rate per 100,000 and 5-yr. count [2003-2007]	17.2 (±4.9) 63	14.6 (±2.6) 129	9.5 (±1.6) 133	20.3 (±4.9) 87	16.0 (±2.9) 122	11.0 (±2.5) 95	15.1 (±2.4) 150	11.8 (±2.1) 119	13.4 (±0.9) 898	14.6 [2006]	MA 6.8 [2006]
Hip Fracture Hospital Discharges Among 65+ Year Olds, rate per 100,000 and 5 yr. count [2004-2008]	704.7 (±65.3) 448	789.2 (±49) 995	738.7 (±39) 1,376	776.3 (±64.1) 563	695.3 (±47.2) 835	746.9 (±49.9) 860	712.8 (±43.9) 1,011	586.3 (±39.9) 829	715.4 (±16.9) 6,917 [2004-2008]	772.0 (±1.3) 1,404,000 [2002-2006]	n/a
Reported Rapes, rate per 10,000 female population and annual average [2003-2007]	3.0 (2.3- 3.9) 11	6.8 (6.0- 7.6) 59	5.4 (4.8- 5.9) 76	1.9 (1.4- 2.6) 8	3.3 (2.7- 3.9) 25	2.2 (1.8- 2.7) 19	7.7 (6.9- 8.5) 76	6.7 (6.1- 7.5) 69	5.1 (4.9- 5.4) 344	n/a	n/a
Traumatic Brain Injury Deaths, age-adjusted rate per 100,000 and 5-yr. count [2003-2007]	21.7 (±5.2) 89	16.1 (±2.6) 147	15.3 (±2.0) 218	24.5 (±4.7) 113	18.5 (±3.0) 155	14.7 (±2.6) 126	17.8 (±2.6) 179	18.8 (±2.7) 194	17.5 (±1.0) 1,221	18.2 [2005]	MA 8.3 [2005]
Unintentional Poisoning Deaths, age-adjusted rate per 100,000 and 5-yr. count [2003-2007]	6.7 (±3.3) 25	11.2 (±2.5) 96	10.2 (±1.7) 139	11.3 (±3.9) 44	10.9 (±2.8) 77	11.4 (±2.6) 93	8.3 (±2.1) 79	9.4 (±2.1) 93	10.0 (±0.8) 646	9.2 [2006]	MD 1.0 [2006]
Firearms Deaths, age-adjusted rate per 100,000 and 5-yr. count [2003-2007]	8.8 (±3.4) 37	7.7 (±2.0) 71	5.6 (±1.4) 80	8.7 (±3.2) 42	7.7 (±2.2) 64	7.4 (±2.1) 64	6.8 (±1.8) 69	7.9 (±1.9) 82	7.3 (±0.6) 509	10.2 151,537 [2002-06]	HI 2.7 [2002-06]
Domestic Assaults Reported to the Police, rate per 10,000 population and count [2007]	31.8 (27.8- 36.2) 229	60.9 (57.2- 64.7) 1,050	40.7 (38.4- 43.2) 1,122	21.2 (18.2- 24.5) 182	29.7 (27.0- 32.6) 447	31.5 (28.8- 34.3) 522	61.8 (58.4- 65.4) 1,196	50.8 (47.7- 54.0) 1,023	43.8 (42.7- 45.0) 5,771	n/a	n/a
Number of substantiated child abuse cases by district of residence (children under age 18), count [2008]	130	402	445	198	196	365	429	352	2,157	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

SUBSTANCE ABUSE AND MENTAL HEALTH INDICATORS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Adults With >=14 Days of Frequent Mental Distress in Past Month, percent [2008]	13.9 (±5.3)	10.7 (±2.7)	10.6 (±2.8)	8.5 (±2.7)	9.9 (±2.2)	10.4 (±2.4)	10.4 (±2.3)	9.7 (±2.6)	10.4 (±1.0)	n/a	n/a
Adults With Current Symptoms of Moderate or Severe Depression, percent [2008]	13.3 (±6.8)	9.0 (±3.2)	7.3 (±2.4)	9.7 (±3.7)	7.2 (±2.1)	7.2 (±3.0)	10.0 (±3.7)	4.5 (±2.0)	8.1 (±1.1)	n/a	n/a
Suicide Deaths, Age >=10 Years, rate per 100,000 [2003-2007]	13.7 (±4.4)	13.2 (±2.5)	11.4 (±1.8)	13.7 (±4.0)	15.7 (±2.9)	12.9 (±2.5)	10.9 (±2.1)	13.5 (±2.3)	12.8 (±.9)	12.8 [2002-2006]	DC 6.3 [2002-06]
Previous 30-Day Alcohol Use, 9 th -12 th Graders, percent [2008]	33.7	32.5	37.8	34.7	38.0	32.5	34.5	34.9	35.0	n/a	n/a
Binge Drinking Within Last 2 Wks, 9 th -12 th Graders, percent [2008]	17.5	17.4	20.4	18.6	20.2	16.4	17.7	16.9	18.2	n/a	n/a
Previous 30-Day Prescription Drug Misuse, 9 th -12 th Graders, percent [2008]	5.4	6.9	7.2	6.2	8.9	6.9	7.2	7.4	7.2	n/a	n/a
Substance Abuse Admissions, rate per 100,000 [2008]	1,527.7	1,425.1	2,559.3	1,570.6	1,308.8	1,707.6	1,123.1	1,019.9	1,593.8	n/a	n/a
Adults Taking Medicine or Receiving Treatment from a Doctor for Mental Health Condition, percent [2008]	15.1 (±6.0)	15.0 (±3.7)	13.4 (±2.9)	16.3 (±5.2)	15.3 (±3.0)	14.7 (±4.1)	17.1 (±3.7)	14.0 (±3.8)	15.0 (±1.2)	n/a	n/a
Adults who Have Ever Had Depression, percent [2008]	26.5 (±8.5)	18.4 (±4.1)	23.1 (±4.3)	17.7 (±4.4)	20.7 (±3.4)	17.9 (±4.3)	23.6 (±4.4)	20.2 (±5.3)	20.9 (±1.7)	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

ACCESS TO CARE INDICATORS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Benchmark State (healthiest)
Access to Primary Care Physician, population to physician ratio [2004]	1,034:1	971:1	759:1	880:1	1,189:1	939:1	1,091:1	1,269:1	978:1	1,351:1 [2000]	MA 187.3 per 100,000 [2005]
Adults With No Health Insurance, percent [2008]	10.8 (±4.7)	15.0 (±4.1)	7.0 (±2.3)	16.8 (±4.2)	14.6 (±3.2)	9.4 (±2.9)	13.9 (±3.3)	6.0 (±2.0)	11.3 (±1.2)	14.5	MA 4.4
Children age 0-18 Years Without Health Insurance, percent and count [2006]	7.8 (7.4-8.2) 1,169	6.3 (6.1-6.6) 2,426	6.2 (6.0-6.4) 3,938	9.1 (8.6-9.5) 1,627	8.0 (7.7-8.3) 2,649	7.4 (7.1-7.6) 2,629	5.4 (5.2-5.6) 2,384	6.6 (6.4-6.9) 3,121	6.8 (6.7-6.9) 19,943	11.7	MA 5.5
PUBLIC HEALTH PREPAREDNESS INDICATORS											
Hospitals with draft hazard plan or a completed hazard plan, percent [2009]	100%	90%	100%	95%	100%	100%	100%	100%	97%	n/a	n/a
Influenza Surveillance: Hospitals Participation in Early Aberration Reporting System (EARS), percent and count [2009]	25.0 1	50.0 2	66.7 2	20.0 1	20.0 1	33.3 2	60.0 3	0 0	34.2 12	n/a	n/a
Influenza Surveillance: Hospitals Participation in Sentinel Provider Program, percent count [2009]	50.0 2	25.0 1	33.3 1	60.0 3	40.0 2	50.0 3	80.0 4	33.3 1	48.6 17	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

PREVENTABLE HOSPITALIZATION COST SAVINGS	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Diabetes short-term complication admission, rate per 100,000 and cost savings if rate declined by 10% [2007]	30.4 \$13,066	51.7 \$53,285	30.9 \$55,963	39.4 \$13,513	50.4 \$30,272	44.6 \$34,944	56.0 \$60,179	33.4 \$32,716	40.8 \$293,939	n/a	n/a
Perforated Appendix admissions, rate per 100,000 patients discharged with appendicitis and cost savings if rate declined by 10% [2007]	51727 \$26,200	48777 \$56,146	31741 \$107,502	40286 \$31,055	43414 \$61,939	32984 \$26,718	28327 \$38,818	33121 \$58,543	30732 \$405,680	n/a	n/a
Diabetes long-term complication admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	128.5 \$78,777	106.0 \$241,354	76.6 \$200,812	80.1 \$60,482	77.7 \$117,343	91.1 \$117,056	93.1 \$137,170	46.6 \$114,148	90.1 1067,142	n/a	n/a
Chronic obstructive pulmonary disease admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	360.2 \$190,225	18.35 \$230,651	151.0 \$206,753	187.7 \$130,817	179.4 \$219,370	268.2 \$230,470	245.6 \$301,067	173.1 \$218,760	224.0 1728,114	n/a	n/a
Hypertension admissions, rate per 100,000 and cost savings if rate declined by 10%, [2007]	38.9 \$15,460	34.0 \$32,799	17.5 \$20,743	18.5 \$8,154	19.6 \$16,367	24.7 \$15,921	13.5 \$10,066	7.6 \$13,262	21.3 \$132,772	n/a	n/a
Congestive heart failure admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	449.8 \$240,959	335.5 \$504,460	319.0 \$522,024	339.0 \$219,780	254.7 \$346,890	375.0 \$361,646	286.1 \$345,249	326.6 \$410,881	352.2 2951,888	n/a	n/a
Low birth weight admissions, rate per 100,000 live births and cost savings if rate declined by 10% [2007]	3439 \$25,077	5913 \$314,790	5145 \$245,139	4489 \$147,590	5408 \$176,735	5553 \$343,157	5188 \$223,391	5484 \$229,725	5264 1705,603	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average

2010 MAINE STATE PROFILE of SELECTED PUBLIC HEALTH INDICATORS

Maine Center for Disease Control and Prevention/DHHS

PREVENTABLE HOSPITALIZATION COST SAVINGS Continued	DISTRICT								COMPARISONS		
	Aroostook (±M.E. or C.I.)	Central (±M.E. or C.I.)	Cumber. (±M.E. or C.I.)	Downeast (±M.E. or C.I.)	Mid Coast (±M.E. or C.I.)	Penquis (±M.E. or C.I.)	Western (±M.E. or C.I.)	York (±M.E. or C.I.)	MAINE STATE (±M.E. or C.I.)	UNITED STATES	Bench- mark State (healthiest)
Dehydration admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	98.1 \$36,885	112.7 \$116,203	71.1 \$70,820	72.7 \$26,942	97.3 \$80,741	118.9 \$84,663	100.3 \$94,081	87.5 \$68,815	101.7 \$579,150	n/a	n/a
Bacterial pneumonia admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	497.2 \$262,085	365.9 \$549,017	235.0 \$363,722	391.4 \$263,525	455.6 \$392,798	468.3 \$454,840	370.9 \$502,517	322.2 \$457,509	379.6 \$324,013	n/a	n/a
Urinary tract infection admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	189.4 \$93,309	105.8 \$121,090	128.3 \$158,612	116.4 \$56,008	135.8 \$113,929	185.3 \$143,702	156.2 \$148,362	120.8 \$105,688	147.7 \$940,703	n/a	n/a
Angina without procedure admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	140.4 \$43,510	47.9 \$37,171	13.8 \$12,996	74.9 \$20,790	32.7 \$22,053	72.4 \$47,184	25.3 \$18,194	15.7 \$11,939	44.7 \$213,837	n/a	n/a
Uncontrolled diabetes admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	8.5 \$2,417	6.5 \$9,325	6.9 \$6,260	14.0 \$6,055	11.1 \$7,716	2.9 \$1,142	6.8 \$3,764	3.8 \$3,264	7.2 \$39,943	n/a	n/a
Adult asthma admissions, rate per 100,000 and cost savings if rate declined by 10% [2007]	108.2 \$49,307	45.8 \$68,817	43.4 \$49,041	45.8 \$26,098	70.7 \$58,999	108.8 \$75,623	72.8 \$64,232	51.6 \$47,515	71.7 \$439,633	n/a	n/a
Lower extremity amputation due to diabetes, rate per 100,000 and cost savings if rate declined by 10% [2007]	28.7 \$28,508	29.8 \$107,479	28.6 \$98,919	27.6 \$30,739	15.3 \$41,829	33.2 \$65,272	21.2 \$58,903	22.7 \$88,076	28.0 \$519,723	n/a	n/a

n/a=not available

Highlighting indicates that the district is statistically different than the state average